

Didymocarpus puhoatensis (Gesneriaceae), a new species from Vietnam

Xin Hong^{1,3}, Zhen-Long Li¹, Stephen Maciejewski², Fang Wen³, Truong Van Do⁴

1 School of Resources and Environmental Engineering, Anhui University, CN-230601, Hefei, China
2 The Gesneriad Society, Inc. 1122 East Pike Street, PMB 637, Seattle, Washington, USA **3** Gesneriad Conservation Centre of China, Guangxi Key Laboratory of Plant Conservation and Restoration Ecology in Karst Terrain, Guangxi Institute of Botany, Guangxi Zhuang Autonomous Region and Chinese Academy of Sciences, CN-541006 Guilin, China **4** Vietnam National Museum of Nature, Vietnam Academy of Science & Technology, 18 Hoang Quoc Viet, Hanoi, Vietnam

Corresponding author: Fang Wen (wenfang760608@139.com); Truong Van Do (dovantruong_bttm@yahoo.com)

Academic editor: K. Ming | Received 17 October 2017 | Accepted 28 November 2017 | Published 29 January 2018

Citation: Hong X, Li Z-L, Maciejewski S, Wen F, Do TV (2017) *Didymocarpus puhoatensis* (Gesneriaceae), a new species from Vietnam. In: Jin X-H, Shui Y-M, Tan Y-H, Kang M (Eds) Plant diversity in Southeast Asia. PhytoKeys 94: 87–93. <https://doi.org/10.3897/phytokeys.94.21650>

Abstract

Didymocarpus puhoatensis, a new species from Vietnam is described and illustrated with photographs. The new species is morphologically similar to *D. brevicalyx* and *D. epithemoides*, but can be easily distinguished by a combination of characters. So far, five species have been recorded in the genus *Didymocarpus* from Vietnam.

Keywords

Didymocarpus, Gesneriaceae, plant taxonomy, Vietnam

Introduction

The delimitation of the genus *Didymocarpus* Wallich (1819: 378) has varied considerably over time (Burtt 1998, Weber et al. 2000, 2011, Möller et al. 2011, Möller and Clark 2013, Li et al. 2015). Now approximately 70 species range from northwest India, eastwards through Nepal, Bhutan, northeast India, Burma (Myanmar), to southern China, Vietnam, Laos, Cambodia, Thailand, the Malay Peninsula and northwards to Sumatra (Weber and Burtt 1998, Weber et al. 2000, Möller et al. 2016). Only three species of this genus were found in Vietnam before 2012, then *D. kerrii* and *D. purpureobracteatus* were respectively reported as new record species for the flora of Vietnam (Phuong et al. 2012,

2014). *Didymocarpus bonii* [= *Calcareoboea bonii*], is now a synonym of *Petrocodon bonii* (Weber et al. 2011). So now there are four species of *Didymocarpus* recorded in Vietnam: *D. kerrii*, *D. pulcher*, *D. poilanei* and *D. purpureobracteatus*.

During a floristic expedition to northern Vietnam in 2015, the authors observed a population of an interesting Gesneriaceae in Pu Hoat Nature Reserve, Nghe An province, Vietnam. It was confirmed that it is member of the genus *Didymocarpus* based on its disc-like stigma (Wang et al. 1998). Over the past two years, the living plants were monitored in the field and an ecological survey was carried out by the co-author in Vietnam and in the nursery of Gesneriads Conservation Centre of China (GCCC) in China.

After thorough comparisons of diagnostic morphological and anatomical features of similar taxa from China, Vietnam, Thailand and adjacent regions (Kiew 1990, Hilliard and Burtt 1995, Wang et al. 1998, Burtt 1998, 1999, Weber et al. 2000, Hilliard 2001, Nangngam and Maxwell 2013, Nangngam and Middleton 2014, Phuong et al. 2014), it is concluded that, as its morphological characters do not fit any known species, it is a new species to science and accordingly described herein. Its morphological characters are compared with the closely related species: *D. brevicalyx* Nangngam & D.J. Middleton (2014: 35) and *D. epithemoides* B.L. Burtt (2001: 92). Therefore, there are five species of the *Didymocarpus* recorded in Vietnam.

Material and methods

Measurements and morphological character assessments of the putative new species were performed and described using specimens' work by the current authors, living material observed in the field and also those cultivated at the Gesneriad Conservation Centre of China. All available specimens of Southeast Asian *Didymocarpus* kept in the following herbaria were examined: E, GH, HN, IBK, K, KUN, MO, PE, PH, US and VNMN. The images of type specimens were also obtained from Tropicos (<http://www.tropicos.org>), JSTOR Global Plants (<http://plants.jstor.org>) and the International Plant Names Index (<http://www.ipni.org>). All morphological characters were studied under dissecting microscopes and are described using the terminology presented by Wang et al. (1990, 1998).

Taxonomic treatment

***Didymocarpus puhoatensis* X.Hong & F.Wen, sp. nov.**

urn:lsid:ipni.org:names:77175491-1

Figures 1, 2

Diagnosis. Although it is morphologically similar to *D. brevicalyx*, it differs by stem densely pubescent, orbicular purple bracts, apices of calyx lobes obtuse, filaments glabrous, staminodes 2; and also similar to *D. epithemoides*, but differs from the latter in having purple calyx, funnel-form corolla, 4–5 cm long, glabrous, dark purple-blackish, ovary glandular puberulent.




Figure 1. *Didymocarpus puhoatensis* X.Hong & F.Wen **A** Habitat **B** Adaxial surface view of leaf blade **C** Adaxial surface view of leaf blade **D** Cyme with flowers, showing the bracts **E** Lateral view of corolla, showing the calyx consisting of a tube.

Type. VIETNAM. Nghệ An Province: Quế Phong, Thông Thụ, Pu Hoat Nature Reserve (Khu Bảo tồn thiên nhiên Pù Hoạt), 19°52'30.5"N, 104°56'15.1"E, alt. 390 m, 18 July 2014, flowering, *Truong Van Do et al.* VNM-CN439 (holotype: IBK; isotype: VNM).

Description. Deciduous, perennial, lithophytic herb, 10–30 cm tall. *Stems* erect, single, sparsely puberulent to glabrescent, the upper, leaf-bearing part and young stems densely covered with whitish multicellular eglandular hairs. *Dry season* plants unknown. *Rainy season* leaves opposite, anisophyllous; petioles terete, 0.5–2.5 cm long, densely covered with multicellular eglandular hairs as on the stems; blades asymmetrically ovate, 6–10 cm long, 5–8 cm wide, apex bluntly acute, base slightly oblique, obtuse-



Figure 2. Flower of *Didymocarpus puhoatensis* X.Hong & F.Wen **A–B** Frontal view of corolla, showing the disc-like stigma **C** Top view of corolla **D** Upward view of corolla **E** Opened corolla, showing stamens and staminodes **F** Pistils without corolla.

cuneate, margin finely serrate or finely doubly serrate, papery, upper surface densely covered with whitish multicellular eglandular hairs, green, lower surface sparsely covered with hairs as on upper surface, pale green, venation pinnate, secondary veins 4–8 on each side of midrib, mostly opposite sometime alternate, obscure above, prominent beneath, covered with whitish multicellular eglandular hairs. *Inflorescences* terminal or from the upper leaf axils, cymose, ca. 12 cm long, 4–10 (–30) flowered; peduncles slender, 6–10 cm long, light green, sparsely covered with multicellular glandular hairs; pedicels 1–1.5 cm long, pale green, with indumentum as on the peduncle. *Bracts* paired, orbicular, ca. 5 mm long and wide, green to pale purple, glabrous. *Calyx* consisting of a tube and shallowly 5-lobed margin, symmetrical, campanulate, 6 mm long, glabrous, somewhat tawny to pinkish purple, tube ca. 5 mm long, 3 mm in diameter;

lobes triangular, ca. 1 mm long, ca. 2 mm wide, apices obtuse. *Corolla* funnellform, 4–5 cm long, glabrous outside, blackish purple, becoming light purple at base; tube ca. 3.5 cm long, base narrow, ca. 2 mm wide, dilated and slightly ventricose towards the throat; widest at throat, diameter ca. 9 mm; corolla bi-lipped, lobes suborbicular; lower lip 3-lobed, ca. 8 mm long, 6 mm wide, more or less equal; upper lip 2-lobed, ca. 4 mm long and wide. Fertile stamens adnate to corolla ca. 1.5 cm from base; anther locules, ca. 2.5 mm long, ca. 1.5 mm wide, densely covered with brownish multicellular eglandular hairs; filaments slender, ca. 1 cm long, white, glabrous; staminodes 3, reduced to thin filaments, lateral ones 3 mm long, the other one 1 mm long, glabrous, adnate to corolla ca. 1 cm from base. *Disc* cylindrical, ca. 2 mm long, margin irregular. *Pistil* ca. 2–3 cm long; ovary narrowly linear, ca. 2 cm long, sparsely glandular puberulent, base reddish, with purple tinge towards stigma and apex green; stigma 1, peltate, concave, papillose, cream. *Capsules* unknown.

Phenology. It flowers and fruits from June to September.

Etymology. The specific epithet is derived from the type locality, Pu Hoat Nature Reserve, Nghệ An Province, Vietnam.

Distribution and habitat. This new taxon is an endemic species from Pu Hoat Nature Reserve of Vietnam. The species grows on limestone rocks in tropical monsoon forest with sufficient seasonal run-off water, at an elevation of 390 m a.s.l. It distributes much lower in altitude and the habitat is much hotter and more humid than other species with stems of the genus.

Preliminary conservation assessment. As population information of *Didymocarpus puhoatensis* is still unclear, it is not appropriate to make an assessment of the extinction risk faced by this new taxon. Thus, the category of Data Deficient (DD) should be appropriate, according to IUCN (2016) criteria. Fortunately, the known habitat of the species is protected as part of a nature reserve. Besides prolonged droughts and illegal logging in the area, there are other potential risks to the persistence of this new species.

Key to the species of *Didymocarpus* in the Vietnam

- 1 Ovary smooth.....2
- Ovary hair4
- 2 Corolla outer hair; leaf blade ovate or elliptic, ca. 4–10 × 2–6 cm.....*D. pulcher*
- Corolla outer smooth; leaf blade near round or broadly elliptic, ca. 1–4 × 1–4 cm*D. kerrii*
- 4 Base of leaf often axisymmetric; bract lanceolata, ca. 2–3 mm long; sepal tube ca. 3 mm long, sepal lobed ca. 1–1.5 mm long*D. poilanei*
- Base of leaf often oblique; bract elliptic ovate or orbicular, ca. 3–8 mm long; sepal tube ca. 8–9 mm long, sepal lobed ca. 2–3 mm long.....5
- 5 Peduncle 4–10 cm, glabrous; bracts connate at base; calyx 1–1.2 cm; corolla 3–4 cm, purple to pinkish purple.....*D. purpureobracteatus*
- peduncles 6–10 cm, glandular hairs; bracts separate; calyx 6 mm; corolla 4–5 cm, blackish purple.....*D. puhoatensis*

Discussion

It is morphologically similar to *D. brevicalyx* and *D. epithemoides* in having the calyx consisting of a tube, similar shape and colour of corolla, both morphological affinities being distributed in Thailand. However, *D. puhoatensis* can be clearly differentiated from both by several characters. The major differences between the species are outlined in Table 1.

Table 1. Diagnostic characters for *Didymocarpus puhoatensis* sp. nov. and its relatives.

Character	<i>Didymocarpus puhoatensis</i>	<i>D. brevicalyx</i>	<i>D. epithemoides</i>
Indumentum of Stem	densely pubescent	densely glandular pubescent	densely pubescent
Bracts	orbicular c. 5 mm long and wide	triangular c. 2 mm long and 1.5 mm wide	orbicular, 5 mm long and wide
Calyx	lobes apices obtuse, purple	lobes apices acute, reddish	lobes apices rounded, violet
Corolla	funnelform, 4–5 cm long, glabrous, dark purple- blackish	funnelform, 4.5 cm long, glabrous, dark purple- blackish	salverform, 3–3.5 cm long, glandular pubescent outside, dark violet
filaments	glabrous	gland-tipped hairs on the upper part	glabrous
Indumentum of ovary	sparsely glandular puberulent	densely glandular pubescent	glabrous

Acknowledgements

This study was financially supported by the Anhui University Doctor Startup Fund, Key University Science Research Project of Anhui Province (No. KJ2017A022), Fund of Guangxi Key Laboratory of Plant Conservation and Restoration Ecology in Karst Terrain (16-B-01-01), Plant germplasm resources projects of the germplasm bank of Wild species of Kunming Institute of Botany, Chinese Academy of Sciences (WGB-1411), the Guangxi Natural Science Foundation (2015GXNSFBB139004 & 2016GXNSFBA380071) and the Key Research and Development Project of Guangxi (Guike AB16380053).

References

Burt BL (1998) Taxonomic history of *Didymocarpus* and *Henckelia* (Gesneriaceae). *Beitrage zur Biologie der Pflanzen*, Breslau 70: 365–375.

Burt BL (1999) Old World Gesneriaceae. VI. Six miscellaneous notes. *Edinburgh Journal of Botany* 56: 371–379. <https://doi.org/10.1017/S0960428600001335>

Burt BL (2001) Flora of Thailand: annotated checklist of Gesneriaceae. *Thai Forest Bulletin (Botany)* 29: 81–109.

Hilliard OM, Burt BL (1995) Old World Gesneriaceae. IV. Notes on *Didymocarpus* and *Lysionotus*. *Edinburgh Journal of Botany* 52: 215–224.

- Hilliard OM (2001) Gesneriaceae. In: Grierson AJC, Long DG, Springate LS (Eds) Flora of Bhutan. Vol 2. Part 3. Royal Botanic Garden, Edinburgh, 1296–1330. <https://doi.org/10.1017/S0960428600000974>
- IUCN (2016) Guidelines for Using the IUCN Red List Categories and Criteria. Version 12. Prepared by the Standards and Petitions Subcommittee. Available from: <http://www.iucn-redlist.org/documents/RedListGuidelines.pdf> [accessed 15 April 2016]
- Kiew R (1990) *Didymocarpus* (Gesneriaceae) on Gunung Tahan, Malaysia. Garden's Bulletin Singapore 42: 47–64.
- Li JM, Sun WJ, Chang Y, Yang WG (2015) Systematic position of *Gyrocheilos* and some odd species of *Didymocarpus* (Gesneriaceae) inferred from molecular data, with reference to pollen and other morphological characters. Journal of Systematics and Evolution 54(2): 113–122. <http://dx.doi.org/10.1111/jse.12169>
- Möller M, Clark JL (2013) The state of molecular studies in the family Gesneriaceae: A review. Selbyana 31(2): 95–125.
- Möller M, Forrest A, Wei YG, Weber A (2011) A molecular phylogenetic assessment of the advanced Asiatic and Malesian didymocarpoid Gesneriaceae with focus on non-monophyletic and monotypic genera. Plant Systematics and Evolution 292: 223–248. <http://dx.doi.org/10.1007/s00606-010-0413-z>
- Möller M, Wei Y-G, Wen F, Clark JL, Weber A (2016) You win some you lose some: updated generic delineations and classification of Gesneriaceae – implications for the family in China. Guihaia 36: 44–60.
- Nangngam P, Maxwell JF (2013) *Didymocarpus* (Gesneriaceae) in Thailand. Garden's Bulletin Singapore 65: 185–225.
- Nangngam P, Middleton DJ (2014) Five new species of *Didymocarpus* (Gesneriaceae) from Thailand. Thai Forest Bulletin (Botany) 42: 35–42.
- Phuong VX, Dang QV, Xuyen DT (2014) Genus *Didymocarpus* Wall. and a new record of species *Didymocarpus purpureobracteatus* Smith for the flora of Vietnam from Xuan Lien natural reserve, Thanh Hoa province. Tap Chi Sinh Hoc 36(1): 45–49. <https://doi.org/10.15625/0866-7160/v36n1.4516>
- Phuong VX, Xuyen DT (2012) A new record of species *Didymocarpus kerrii* Craib (Gesneriaceae Dumort) for the flora of Vietnam. Tap Chi Sinh Hoc 34(1): 404–405.
- Wallich N (1829) A numerical list of dried specimens of plants in the East India Company's Museum: collected under the superintendence of Dr. Wallich of the Company's botanic garden at Calcutta. East India Company, London, 306 pp.
- Wang WT, Pan KY, Li ZU, Weitzman AL, Skog LE (1998) Gesneriaceae. In: Wu ZY, Raven PH (Eds) Flora of China. Vol. 18. Science Press, Beijing, 349–358.
- Weber A, Burtt BL (1998) Remodelling of *Didymocarpus* and associated genera (Gesneriaceae). Beiträge zur Biologie der Pflanzen 70: 293–363.
- Weber A, Burtt VL, Vitek E (2000) Materials for a revision of *Didymocarpus* (Gesneriaceae). Annalen des Naturhistorischen Museums in Wien 102: 441–475.
- Weber A, Wei YG, Puglisi C, Wen F, Mayer V, Möller M (2011) A new definition of the genus *Petrocodon* (Gesneriaceae). Phytotaxa 23(1): 49–67. <http://dx.doi.org/10.11646/phyto-taxa.23.1.3>